

Advanced Counting Techniques

Generating Functions

Abhijit Das

Department of Computer Science and Engineering
Indian Institute of Technology Kharagpur

October 1, 2020

A Counting Problem

You appear in four tests.

- Algorithms
- Bioinformatics
- Compilers
- Discrete Structures

In each test, you get an integer mark in the range $[0, 10]$.

In how many ways can you get a total of 25 marks?

Some examples: $5 + 5 + 10 + 5 = 10 + 5 + 5 + 5 = 6 + 7 + 6 + 6 = 1 + 9 + 8 + 7 = 25$.

Frame the Problem Algebraically

- Algorithms: $A = 1 + a + a^2 + a^3 + a^4 + a^5 + a^6 + a^7 + a^8 + a^9 + a^{10}$.
- Bioinformatics: $B = 1 + b + b^2 + b^3 + b^4 + b^5 + b^6 + b^7 + b^8 + b^9 + b^{10}$.
- Compilers: $C = 1 + c + c^2 + c^3 + c^4 + c^5 + c^6 + c^7 + c^8 + c^9 + c^{10}$.
- Discrete Structures: $D = 1 + d + d^2 + d^3 + d^4 + d^5 + d^6 + d^7 + d^8 + d^9 + d^{10}$.

Consider the product $ABCD$.

The answer is the number of terms of the form $a^i b^j c^k d^l$ in $ABCD$ with $i + j + k + l = 25$.

No real progress actually.

An Insight

We are considering terms $a^i b^j c^k d^l$ with $i + j + k + l = 25$.

We can take $a = b = c = d = x$.

The coefficient of x^{25} in

$$\begin{aligned}(1 + x + x^2 + x^3 + \cdots + x^{10})^4 &= \left(\frac{1 - x^{11}}{1 - x} \right)^4 \\ &= \left(1 - \binom{4}{1} x^{11} + \binom{4}{2} x^{22} - \binom{4}{3} x^{33} + x^{44} \right) \sum_{i \geq 0} \binom{i+3}{i} x^i\end{aligned}$$

gives the answer

$$\binom{25+3}{25} - \binom{4}{1} \binom{14+3}{14} + \binom{4}{2} \binom{3+3}{3} = \binom{28}{25} - \binom{4}{1} \binom{17}{14} + \binom{4}{2} \binom{6}{3} = 676.$$

Exercise: Deduce the same formula by the principle of inclusion and exclusion.

Combination with Repetitions

To choose r objects with repetition from a set of n distinct objects.

Each object can be chosen a maximum of r times.

Look at the coefficient of x^r in $(1 + x + x^2 + \dots + x^r)^n$.

To simplify matters, look at the infinite series

$$\begin{aligned}(1 + x + x^2 + \dots)^n &= \left(\frac{1}{1-x}\right)^n \\ &= \frac{1}{(1-x)^n} \\ &= \sum_{i \geq 0} \binom{n+i-1}{i} x^i.\end{aligned}$$

The coefficient of x^r is $\binom{n+r-1}{r}$.

Definition

Let $a_0, a_1, a_2, a_3, \dots, a_n, \dots$ be an infinite sequence of real numbers.

The **generating function** of the sequence is

$$A(x) = a_0 + a_1x + a_2x^2 + \dots + a_nx^n + \dots$$

The power series $A(x)$ is *formal*.

We usually do not put any value for x in $A(x)$.

Consequently, the convergence of the series is usually not an issue.

If we want to put a value for x , convergence issues must be considered.

Examples

- Let $n \in \mathbb{N}$. Then $(1+x)^n$ is the generating function of

$$\binom{n}{0}, \binom{n}{1}, \binom{n}{2}, \dots, \binom{n}{n}, 0, 0, 0, \dots$$

- Let $n \in \mathbb{N}$. Then $\frac{1-x^n}{1-x} = 1+x+x^2+\dots+x^{n-1}$ is the generating function of

$$\underbrace{1, 1, 1, \dots, 1}_{n \text{ times}}, 0, 0, 0, \dots$$

- $\frac{1}{1-x} = 1+x+x^2+\dots$ is the generating function of $1, 1, 1, \dots$

Examples

- $\frac{1}{(1-x)^2} = 1 + 2x + 3x^2 + 4x^3 + \dots + (n+1)x^n + \dots$

is the generating function of $1, 2, 3, 4, 5, \dots$

- $\frac{x}{(1-x)^2} = 0 + x + 2x^2 + 3x^3 + 4x^4 + \dots + nx^n + \dots$

is the generating function of $0, 1, 2, 3, 4, 5, \dots$

- $\frac{d}{dx} \frac{x}{(1-x)^2} = \frac{1+x}{(1-x)^3} = 1^2 + 2^2x + 3^2x^2 + 4^2x^3 + \dots + (n+1)^2x^n + \dots$

is the generating function of $1^2, 2^2, 3^2, 4^2, 5^2, \dots$

- $\frac{x(1+x)}{(1-x)^3}$ is the generating function of $0^2, 1^2, 2^2, 3^2, 4^2, 5^2, \dots$

Examples

- $\frac{1}{1-\alpha x} = 1 + \alpha x + \alpha^2 x^2 + \alpha^3 x^3 + \cdots + \alpha^n x^n + \cdots$

is the generating function of the geometric series $1, \alpha, \alpha^2, \alpha^3, \dots, \alpha^n, \dots$

- If $A(x)$ is the generating function of $a_0, a_1, a_2, \dots, a_n, \dots$, and $B(x)$ the generating function of $b_0, b_1, b_2, \dots, b_n, \dots$, then $A(x) + B(x)$ is the generating function of $a_0 + b_0, a_1 + b_1, a_2 + b_2, \dots, a_n + b_n, \dots$

- $A(x)B(x)$ is the generating function of the **convolution**
 $a_0b_0, a_0b_1 + a_1b_0, a_0b_2 + a_1b_1 + a_2b_0, \dots, a_0b_n + a_1b_{n-1} + a_2b_{n-2} + \cdots + a_nb_0, \dots$

- Take $B(x) = \frac{1}{1-x}$ in the convolution to see that $\frac{A(x)}{1-x}$ is the generating function of the **prefix sums** $a_0, a_0 + a_1, a_0 + a_1 + a_2, \dots, a_0 + a_1 + a_2 + \cdots + a_n, \dots$

Examples

In how many ways 20 marbles can be placed in three boxes such that

- (1) Each box contains at least two marbles, and
- (2) The third box contains no more than ten marbles?

Look at the coefficient of x^{20} in

$$\begin{aligned} & (x^2 + x^3 + x^4 + \cdots)^2 (x^2 + x^3 + \cdots + x^{10}) \\ &= x^6 (1 + x + x^2 + \cdots)^2 (1 + x + x^2 + \cdots + x^8) \\ &= \frac{x^6 (1 - x^9)}{(1 - x)^3} \\ &= (x^6 - x^{15}) \sum_{i \geq 0} \binom{i+2}{i} x^i. \end{aligned}$$

The answer is $\binom{14+2}{14} - \binom{5+2}{5} = \binom{16}{2} - \binom{7}{2} = 120 - 21 = 99$.

Examples

How many 5-element subsets of $\{1, 2, 3, 4, \dots, 20\}$ do not contain consecutive integers?

Let $\{a_1, a_2, a_3, a_4, a_5\}$ be such a subset with

$$1 = a_0 \leq a_1 < a_2 < a_3 < a_4 < a_5 \leq a_6 = 20.$$

For $i = 0, 1, 2, 3, 4, 5$, define $d_i = a_{i+1} - a_i$.

We have $d_0, d_5 \geq 0$, $d_1, d_2, d_3, d_4 \geq 2$, and $d_0 + d_1 + d_2 + d_3 + d_4 + d_5 = 20 - 1 = 19$.

The answer is the coefficient of x^{19} in

$$\begin{aligned} & (1 + x + x^2 + \dots)^2 (x^2 + x^3 + x^4 + \dots)^4 \\ &= \frac{x^8}{(1-x)^6} = x^8 \sum_{i \geq 0} \binom{i+5}{i} x^i, \end{aligned}$$

that is, $\binom{11+5}{11} = \binom{16}{5} = 4368$.

Geometric Distribution

- You toss a coin repeatedly until a head occurs.
- In each toss, p is the probability of head.
- Probability of tail is $q = 1 - p$ in each toss.
- Assume that $0 < p < 1$, so $0 < q < 1$ too.
- Let G be the number of times you need to toss.
- G assumes positive integral values.
- $\Pr[G = n] = q^{n-1}p$ for $n = 1, 2, 3, \dots$
- We want to compute $E[G]$ and $\text{Var}[G]$.

Expectation

- We have $\frac{1}{(1-x)^2} = 1 + 2x + 3x^2 + 4x^3 + \dots + nx^{n-1} + \dots$.
- The series converges for $|x| < 1$.
- Put $x = q$ to get

$$\frac{1}{(1-q)^2} = \frac{1}{p^2} = 1 + 2q + 3q^2 + 4q^3 + \dots + nq^{n-1} + \dots$$

- $E[G] = p + 2qp + 3q^2p + 4q^3p + \dots + nq^{n-1}p + \dots = p \times \frac{1}{p^2} = \frac{1}{p}$.

Variance

- $\text{Var}(G) = E[G^2] - E[G]^2$.
- We have seen that $\frac{1+x}{(1-x)^3} = 1^2 + 2^2x + 3^2x^2 + \dots + n^2x^{n-1} + \dots$.
- This series too converges for $|x| < 1$.
- Put $x = q$ to get

$$1^2 + 2^2q + 3^2q^2 + \dots + n^2q^{n-1} + \dots = \frac{1+q}{(1-q)^3} = \frac{1+q}{p^3}.$$

- $E[G^2] = 1^2p + 2^2qp + 3^2q^2p + 4^2q^3p + \dots + n^2q^{n-1}p + \dots = p \times \left(\frac{1+q}{p^3}\right) = \frac{1+q}{p^2}$.
- Thus $\text{Var}(G) = \frac{1+q}{p^2} - \left(\frac{1}{p}\right)^2 = \frac{q}{p^2}$.

Compositions and Partitions of Positive Integers

Abhijit Das

Department of Computer Science and Engineering
Indian Institute of Technology Kharagpur

October 1, 2020

Ordered and Unordered Partitions of Positive Integers

Let $n \in \mathbb{N}$.

In how many ways can n be written as a sum of positive integers?

If the order of the summands is important, we talk about **compositions**.

If the order of the summands is not important, we talk about **partitions**.

Compositions of 4 are

$$4 = 1 + 3 = 3 + 1 = 2 + 2 = 1 + 1 + 2 = 1 + 2 + 1 = 1 + 1 + 2 = 1 + 1 + 1 + 1.$$

Partitions of 4 are $4 = 1 + 3 = 2 + 2 = 1 + 1 + 2 = 1 + 1 + 1 + 1$.

We proved earlier that the number of compositions of n is 2^{n-1} .

The number of partitions of n does not have a known closed-form formula.

We will study these again in the light of generating functions.

Counting Compositions of n

- Classify compositions by number of summands.
- One summand: Only one way of writing each $n \geq 1$. So the generating function is

$$x + x^2 + x^3 + \cdots + x^n + \cdots = \frac{x}{1-x}.$$

- Two summands: Look at the coefficient of x^n in

$$(x + x^2 + x^3 + \cdots)^2 = \left(\frac{x}{1-x}\right)^2.$$

- In general, for i summands, consider the coefficient of x^n in

$$(x + x^2 + x^3 + \cdots)^i = \left(\frac{x}{1-x}\right)^i.$$

Counting Compositions of n

The generating function of the number of compositions of n is

$$\begin{aligned}\sum_{i \geq 1} \left(\frac{x}{1-x}\right)^i &= \left(\frac{x}{1-x}\right) \sum_{i \geq 0} \left(\frac{x}{1-x}\right)^i \\ &= \left(\frac{x}{1-x}\right) \left[\frac{1}{1 - \left(\frac{x}{1-x}\right)} \right] \\ &= \frac{x}{1-2x} \\ &= x(1 + 2x + 2^2x^2 + 2^3x^3 + \dots + 2^{n-1}x^{n-1} + \dots) \\ &= x + 2x^2 + 2^2x^3 + 2^3x^4 + \dots + 2^{n-1}x^n + \dots.\end{aligned}$$

We have again derived that the number of compositions of n is 2^{n-1} .

Counting Palindromic Compositions of n

- $4 = 2 + 2 = 1 + 2 + 1 = 1 + 1 + 1 + 1$.
- $5 = 2 + 1 + 2 = 1 + 3 + 1 = 1 + 1 + 1 + 1 + 1$.
- If the number of summands is even, n must be even.
- If the number of summands is odd, then the middle summand must have the same parity as n .
- To the left of the center, any arbitrary composition is possible.
- To the right of the center, we write this composition in the reverse order.

Case 1: n is Odd

- n may be the only summand (one case).
- Now consider multiple summands.
- The number of summands must be odd.
- The central summand must be odd (any one of $1, 3, 5, 7, \dots, n-2$).
- The remaining sum is $n-1, n-3, n-5, n-7, \dots, 2$.
- This is distributed equally to the two sides of the center.
- The total count of palindromic compositions is therefore

$$1 + \left(2^{\frac{n-1}{2}-1} + 2^{\frac{n-3}{2}-1} + 2^{\frac{n-5}{2}-1} + \dots + 2^{1-1} \right) = 2^{\frac{n-1}{2}} = 2^{\lfloor \frac{n}{2} \rfloor}.$$

Case 2: n is Even

- n may be the only summand (one case).
- First, consider odd number of summands.
- The central summand must be even (any one of $2, 4, 6, 8, \dots, n - 2$).
- The remaining sum is $n - 2, n - 4, n - 6, n - 8, \dots, 2$.
- This is distributed equally to the two sides of the center.
- The total count of palindromic compositions with odd number of summands is $1 + \left(2^{\frac{n-2}{2}-1} + 2^{\frac{n-4}{2}-1} + 2^{\frac{n-6}{2}-1} + \dots + 2^{1-1}\right) = 2^{\frac{n-2}{2}} = 2^{\frac{n}{2}-1}$.
- If the number of summands is even, any composition of $n/2$ gives a palindromic composition of n . The count in this case is $2^{\frac{n}{2}-1}$.
- The total count is $2^{\frac{n}{2}} = 2^{\lfloor \frac{n}{2} \rfloor}$.

Counting Partitions of n

- $p(n)$ is the number of partitions of n .
- We need to count how many times each $i \in \mathbb{N}$ may occur in the sum.
- 1 may occur never or once or twice or thrice or ... giving the power series
$$1 + x + x^2 + x^3 + \dots = \frac{1}{1-x}.$$
- 2 may occur never or once or twice or thrice or ... giving the power series
$$1 + x^2 + x^4 + x^6 + \dots = \frac{1}{1-x^2}.$$
- In general, i may occur never or once or twice or thrice or ... giving the power series
$$1 + x^i + x^{2i} + x^{3i} + \dots = \frac{1}{1-x^i}.$$

Counting Partitions of n

- The generating function for $p(n)$ is

$$\prod_{i \geq 1} \frac{1}{1 - x^i}.$$

- We may truncate the product after $i = n$.
- Nevertheless, we do not get any closed-form formula for $p(n)$.

Counting Partitions of n with Distinct Summands

- $p_d(n)$ is the number of partitions of n with distinct summands.
- $7 = 1 + 6 = 2 + 5 = 3 + 4 = 1 + 2 + 4$.
- $p_d(7) = 5$.
- Each $i \in \mathbb{N}$ occurs never or once in the sum.
- For any $i \geq 1$, the relevant series is $1 + x^i$.
- The generating function for $p_d(n)$ is therefore

$$(1 + x)(1 + x^2)(1 + x^3) \cdots = \prod_{i \geq 1} (1 + x^i).$$

Counting Partitions of n with Odd Summands

- $p_o(n)$ is the number of partitions of n with odd summands.
- $7 = 1 + 1 + 5 = 1 + 3 + 3 = 1 + 1 + 1 + 1 + 3 = 1 + 1 + 1 + 1 + 1 + 1 + 1$.
- $p_o(7) = 5$.
- Only $1, 3, 5, 7, \dots$ are allowed in the sum.
- The generating function for $p_o(n)$ is therefore

$$\begin{aligned} & (1 + x + x^2 + x^3 + \dots)(1 + x^3 + x^6 + x^9 + \dots)(1 + x^5 + x^{10} + x^{15} + \dots) \dots \\ &= \left(\frac{1}{1-x} \right) \left(\frac{1}{1-x^3} \right) \left(\frac{1}{1-x^5} \right) \dots \\ &= \prod_{i \geq 1} \left(\frac{1}{1-x^{2i-1}} \right). \end{aligned}$$



Equality of the Last Two Generating Functions

$$\text{Use } 1 + x^i = \frac{1 - x^{2i}}{1 - x^i}.$$

$$\begin{aligned} \prod_{i \geq 1} (1 + x^i) &= \prod_{i \geq 1} \frac{1 - x^{2i}}{1 - x^i} \\ &= \left(\frac{1 - x^2}{1 - x} \right) \left(\frac{1 - x^4}{1 - x^2} \right) \left(\frac{1 - x^6}{1 - x^3} \right) \left(\frac{1 - x^8}{1 - x^4} \right) \cdots \\ &= \left(\frac{1}{1 - x} \right) \left(\frac{1}{1 - x^3} \right) \left(\frac{1}{1 - x^5} \right) \cdots \end{aligned}$$

It follows that $p_d(n) = p_o(n)$ for all $n \in \mathbb{N}$.

Ferrers Diagrams



$$20 = 7 + 5 + 4 + 2 + 2 = 5 + 5 + 3 + 3 + 2 + 1 + 1.$$

Observation: The number of partitions of n into m summands is equal to the number of partitions of n into summands, among which m is the largest.

Exercise: Prove that the number of partitions of n is equal to the number of partitions of $2n$ into n summands.

Exponential Generating Functions

$$\begin{aligned}(1+x)^n &= C(n,0) + C(n,1)x + C(n,2)x^2 + \cdots + C(n,i)x^i + \cdots + C(n,n)x^n \\ &= \frac{P(n,0)}{0!} + \frac{P(n,1)}{1!}x + \frac{P(n,2)}{2!}x^2 + \cdots + \frac{P(n,i)}{i!}x^i + \cdots + \frac{P(n,n)}{n!}x^n.\end{aligned}$$

Let $a_0, a_1, a_2, \dots, a_n, \dots$ be a sequence of real numbers.

The **ordinary generating function** (OGF) of the sequence is

$$A(x) = a_0 + a_1x + a_2x^2 + \cdots + a_nx^n + \cdots.$$

The **exponential generating function** (EGF) of the sequence is

$$A_e(x) = \frac{a_0}{0!} + \frac{a_1}{1!}x + \frac{a_2}{2!}x^2 + \cdots + \frac{a_n}{n!}x^n + \cdots.$$

Exponential generating functions are used for counting arrangements.

Examples

- $e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \cdots + \frac{x^n}{n!} + \cdots$ is the EGF of $1, 1, 1, 1, 1, \dots$, and the OGF of $\frac{1}{0!}, \frac{1}{1!}, \frac{1}{2!}, \frac{1}{3!}, \dots, \frac{1}{n!}, \dots$
- $\frac{e^x + e^{-x}}{2} = 1 + \frac{x^2}{2!} + \frac{x^4}{4!} + \frac{x^6}{6!} + \cdots$ is the EGF of $1, 0, 1, 0, 1, 0, \dots$
- $\frac{e^x - e^{-x}}{2} = x + \frac{x^3}{3!} + \frac{x^5}{5!} + \frac{x^7}{7!} + \cdots$ is the EGF of $0, 1, 0, 1, 0, 1, \dots$

Application 1

How many arrangements of four letters are possible using the letters of ANANAS?

There are three A's, two N's, and one S. So the EGF is

$$\left(1 + x + \frac{x^2}{2!} + \frac{x^3}{3!}\right) \left(1 + x + \frac{x^2}{2!}\right) (1 + x).$$

The term involving x^4 is

$$\underbrace{\left(\frac{x^3}{3!} \times x \times 1\right)}_{\text{AAAN}} + \underbrace{\left(\frac{x^3}{3!} \times 1 \times x\right)}_{\text{AAAS}} + \underbrace{\left(\frac{x^2}{2!} \times \frac{x^2}{2!} \times 1\right)}_{\text{AANN}} + \underbrace{\left(\frac{x^2}{2!} \times x \times x\right)}_{\text{AANS}} + \underbrace{\left(x \times \frac{x^2}{2!} \times x\right)}_{\text{ANNS}}$$

The coefficient of $\frac{x^4}{4!}$ in the EGF is $\frac{4!}{3!} + \frac{4!}{3!} + \frac{4!}{2!2!} + \frac{4!}{2!} + \frac{4!}{2!}$.

Application 2

How many strings of length $n > 0$ over $\{a, b, c, d\}$ such that

- (1) the number of c 's is odd, and
- (2) the number of d 's is even?

The EGF is

$$\begin{aligned} & \left(1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \cdots\right)^2 \left(x + \frac{x^3}{3!} + \frac{x^5}{5!} + \cdots\right) \left(1 + \frac{x^2}{2!} + \frac{x^4}{4!} + \cdots\right) \\ &= e^{2x} \left(\frac{e^x - e^{-x}}{2}\right) \left(\frac{e^x + e^{-x}}{2}\right) = \frac{1}{4} e^{2x} (e^{2x} - e^{-2x}) = \frac{1}{4} (e^{4x} - 1) = \frac{1}{4} \sum_{n \geq 1} \frac{(4x)^n}{n!}. \end{aligned}$$

The coefficient of $\frac{x^n}{n!}$ is 4^{n-1} .

Using Generating Functions to Solve Recurrence Relations

Abhijit Das

Department of Computer Science and Engineering
Indian Institute of Technology Kharagpur

October 1, 2020

A Simple Recurrence

$$a_0 = 1$$

$$a_n = 3a_{n-1} + 2n \text{ for } n \geq 1.$$

The first few terms of the sequence are

$$a_0 = 1,$$

$$a_1 = 3 \times 1 + 2 \times 1 = 5,$$

$$a_2 = 3 \times 5 + 2 \times 2 = 19,$$

$$a_3 = 3 \times 19 + 2 \times 3 = 63,$$

$$a_4 = 3 \times 63 + 2 \times 4 = 197,$$

...

The Generating Function of the Sequence

$$\begin{aligned}A(x) &= a_0 + a_1x + a_2x^2 + a_3x^3 + \cdots + a_nx^n + \cdots \\&= 1 + (3a_0 + 2 \times 1)x + (3a_1 + 2 \times 2)x^2 + (3a_2 + 2 \times 3)x^3 + \cdots + (3a_{n-1} + 2n)x^n + \cdots \\&= 1 + 3x(a_0 + a_1x + a_2x^2 + a_3x^3 + \cdots + a_nx^n + \cdots) + \\&\quad 2x(1 + 2x + 3x^2 + \cdots + nx^{n-1} + \cdots) \\&= 1 + 3xA(x) + \frac{2x}{(1-x)^2}.\end{aligned}$$

Therefore

$$A(x) = \frac{1}{1-3x} + \frac{2x}{(1-x)^2(1-3x)}.$$

Expand $A(x)$

$$A(x) = \frac{1}{1-3x} + \frac{2x}{(1-x)^2(1-3x)} = \frac{A}{1-3x} + \frac{B}{1-x} + \frac{C}{(1-x)^2}.$$

$$(1-x)^2 + 2x = A(1-x)^2 + B(1-x)(1-3x) + C(1-3x).$$

Put $x = \frac{1}{3}$ to get $A = \frac{(1 - \frac{1}{3})^2 + \frac{2}{3}}{(1 - \frac{1}{3})^2} = \frac{4+6}{4} = \frac{5}{2}$.

Put $x = 1$ to get $C = \frac{(1-1)^2 + 2}{1-3} = -1$.

Equate the constant coefficient to get $1 = A + B + C$, so $B = 1 - A - C = 1 - \frac{5}{2} + 1 = -\frac{1}{2}$.

Power Series Expansion of $A(x)$

$$\begin{aligned}A(x) &= \frac{\frac{5}{2}}{1-3x} - \frac{\frac{1}{2}}{1-x} - \frac{1}{(1-x)^2} \\&= \frac{5}{2} \left[1 + 3x + 3^2x^2 + 3^3x^3 + \cdots + 3^n x^n + \cdots \right] \\&\quad - \frac{1}{2} \left[1 + x + x^2 + x^3 + \cdots + x^n + \cdots \right] \\&\quad - \left[1 + 2x + 3x^2 + 4x^3 + \cdots + (n+1)x^n + \cdots \right].\end{aligned}$$

Therefore $a_n = \frac{5}{2} \times 3^n - \frac{1}{2} - (n+1) = \frac{5}{2} \times 3^n - n - \frac{3}{2}$ for all $n \geq 0$.

Fibonacci Numbers

$$F_0 = 0,$$

$$F_1 = 1,$$

$$F_n = F_{n-1} + F_{n-2} \text{ for } n \geq 2.$$

The generating function of the Fibonacci sequence is

$$F(x) = F_0 + F_1x + F_2x^2 + F_3x^3 + \cdots + F_nx^n + \cdots .$$

Derivation of $F(x)$

$$\begin{aligned}F(x) &= F_0 + F_1x + F_2x^2 + F_3x^3 + \cdots + F_nx^n + \cdots \\&= 0 + x + (F_1 + F_0)x^2 + (F_2 + F_1)x^3 + (F_3 + F_2)x^4 + \cdots + (F_{n-1} + F_{n-2})x^n + \cdots \\&= x + (F_1x^2 + F_2x^3 + F_3x^4 + \cdots + F_{n-1}x^n + \cdots) + \\&\quad (F_0x^2 + F_1x^3 + F_2x^4 + \cdots + F_{n-2}x^n + \cdots) \\&= x - F_0x + (F_0x + F_1x^2 + F_2x^3 + F_3x^4 + \cdots + F_{n-1}x^n + \cdots) + \\&\quad (F_0x^2 + F_1x^3 + F_2x^4 + \cdots + F_{n-2}x^n + \cdots) \\&= x + x(F_0 + F_1x + F_2x^2 + F_3x^3 + \cdots + F_nx^n + \cdots) + \\&\quad x^2(F_0 + F_1x + F_2x^2 + F_3x^3 + \cdots + F_nx^n + \cdots) \\&= x + xF(x) + x^2F(x).\end{aligned}$$

Therefore

$$F(x) = \frac{x}{1 - x - x^2}.$$

Playing with $F(x)$

We want to write the denominator as

$$1 - x - x^2 = (1 - \alpha x)(1 - \beta x).$$

Therefore we solve the quadratic equation $y^2 - y - 1 = 0$.

The roots are $\alpha = \frac{1 + \sqrt{5}}{2}$ and $\beta = \frac{1 - \sqrt{5}}{2}$.

This gives $F(x) = \frac{x}{1 - x - x^2} = \frac{x}{(1 - \alpha x)(1 - \beta x)} = \frac{A}{1 - \alpha x} + \frac{B}{1 - \beta x}$.

It follows that $x = A(1 - \beta x) + B(1 - \alpha x)$.

Put $x = 1/\alpha$ and $x = 1/\beta$ to get

$$A = \frac{1/\alpha}{1 - \beta/\alpha} = \frac{1}{\alpha - \beta} = \frac{1}{\sqrt{5}} \quad \text{and} \quad B = \frac{1/\beta}{1 - \alpha/\beta} = \frac{1}{\beta - \alpha} = -\frac{1}{\sqrt{5}}.$$

Explicit Formula for Fibonacci Numbers

We have derived

$$F(x) = \frac{1}{\sqrt{5}} \left[\frac{1}{1 - \alpha x} - \frac{1}{1 - \beta x} \right].$$

Power-series expansions of the two terms give

$$\begin{aligned} F_n &= \frac{1}{\sqrt{5}} (\alpha^n - \beta^n) \\ &= \frac{1}{\sqrt{5}} \left[\left(\frac{1 + \sqrt{5}}{2} \right)^n - \left(\frac{1 - \sqrt{5}}{2} \right)^n \right] \end{aligned}$$

for all $n \geq 0$.

A More Complicated Recurrence

$$b_0 = 1,$$

$$b_1 = 2,$$

$$b_2 = 3,$$

$$b_n = b_{n-1} + b_{n-2} - b_{n-3} + 4 \text{ for all } n \geq 3.$$

A few other terms of the sequence are

$$b_3 = 3 + 2 - 1 + 4 = 8,$$

$$b_4 = 8 + 3 - 2 + 4 = 13,$$

$$b_5 = 13 + 8 - 3 + 4 = 22,$$

$$b_6 = 22 + 13 - 8 + 4 = 31,$$

...

The Generating Function of the Sequence

$$\begin{aligned} B(x) &= b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + \cdots + b_nx^n + \cdots \\ &= 1 + 2x + 3x^2 + \sum_{n \geq 3} (b_{n-1} + b_{n-2} - b_{n-3} + 4)x^n \\ &= 1 + 2x + 3x^2 + x \sum_{n \geq 3} b_{n-1}x^{n-1} + x^2 \sum_{n \geq 3} b_{n-2}x^{n-2} - x^3 \sum_{n \geq 3} b_{n-3}x^{n-3} + 4x^3 \sum_{n \geq 3} x^{n-3} \\ &= 1 + 2x + 3x^2 + x(B(x) - b_0 - b_1x) + x^2(B(x) - b_0) - x^3B(x) + \frac{4x^3}{1-x} \\ &= 1 + 2x + 3x^2 - x - 2x^2 - x^2 + (x + x^2 - x^3)B(x) + \frac{4x^3}{1-x} \\ &= 1 + x + (x + x^2 - x^3)B(x) + \frac{4x^3}{1-x}. \end{aligned}$$

$$\text{Therefore } B(x) = \frac{1 - x^2 + 4x^3}{(1-x)(1-x-x^2+x^3)} = \frac{1 - x^2 + 4x^3}{(1-x)^3(1+x)}.$$

Expand $B(x)$

$$B(x) = \frac{1 - x^2 + 4x^3}{(1-x)^3(1+x)} = \frac{A}{1-x} + \frac{B}{(1-x)^2} + \frac{C}{(1-x)^3} + \frac{D}{1+x}.$$

$$1 - x^2 + 4x^3 = A(1-x)^2(1+x) + B(1-x)(1+x) + C(1+x) + D(1-x)^3.$$

Put $x = 1$ to get $C = 2$.

Put $x = -1$ to get $D = -\frac{1}{2}$.

Coefficient of x^3 : $4 = A - D$, so $A = 4 + D = \frac{7}{2}$.

Constant term: $1 = A + B + C + D$, so $B = 1 - A - C - D = -4$.

Power Series Expansion of $B(x)$

$$\begin{aligned} B(x) &= \frac{\frac{7}{2}}{1-x} - \frac{4}{(1-x)^2} + \frac{2}{(1-x)^3} - \frac{\frac{1}{2}}{1+x} \\ &= \frac{7}{2} \sum_{n \geq 0} x^n - 4 \sum_{n \geq 0} (n+1)x^n + 2 \sum_{n \geq 0} \frac{(n+1)(n+2)}{2!} x^n - \frac{1}{2} \sum_{n \geq 0} (-1)^n x^n. \end{aligned}$$

Therefore

$$\begin{aligned} b_n &= \frac{7}{2} - 4(n+1) + (n+1)(n+2) - \frac{1}{2}(-1)^n \\ &= n^2 - n + \frac{3}{2} - \frac{1}{2}(-1)^n. \end{aligned}$$

Catalan Numbers

$$C_0 = 1,$$

$$C_n = C_0C_{n-1} + C_1C_{n-2} + C_2C_{n-3} + \cdots + C_{n-1}C_0 \text{ for all } n \geq 1.$$

The generating function for the Catalan series is

$$\begin{aligned} C(x) &= C_0 + C_1x + C_2x^2 + C_3x^3 + \cdots + C_nx^n + \cdots \\ &= 1 + C_0C_0x + (C_0C_1 + C_1C_0)x^2 + (C_0C_2 + C_1C_1 + C_2C_0)x^3 + \cdots + \\ &\quad (C_0C_{n-1} + C_1C_{n-2} + \cdots + C_{n-1}C_0)x^n + \cdots \\ &= 1 + x(C_0 + C_1x + C_2x^2 + \cdots + C_nx^n + \cdots)(C_0 + C_1x + C_2x^2 + \cdots + C_nx^n + \cdots) \\ &= 1 + xC(x)^2. \end{aligned}$$

Solve for $C(x)$

$$xC(x)^2 - C(x) + 1 = 0.$$

$$C(x) = \frac{1 \pm \sqrt{1-4x}}{2x}.$$

The choice of $+$ in the numerator gives a term $1/x$ in $C(x)$. So

$$C(x) = \frac{1 - \sqrt{1-4x}}{2x}.$$

Therefore C_n is equal to half the coefficient of x^{n+1} in the power-series expansion of the numerator $1 - \sqrt{1-4x}$.

The Closed-Form Formula for Catalan Numbers

$$\begin{aligned}C_n &= -(-4)^{n+1} \frac{1}{2} \left[\frac{\frac{1}{2}(\frac{1}{2}-1)(\frac{1}{2}-2)(\frac{1}{2}-3)\cdots(\frac{1}{2}-n)}{(n+1)!} \right] \\&= (-1)^{n+2} \times \frac{4^{n+1}}{2^{n+2}} \times (-1)^n \times \left[\frac{1 \times 3 \times 5 \times \cdots \times (2n-1)}{(n+1)!} \right] \\&= 2^n \times \left[\frac{1 \times 3 \times 5 \times \cdots \times (2n-1)}{(n+1)!} \right] \\&= 2^n \times \left[\frac{1 \times 3 \times 5 \times \cdots \times (2n-1) \times n!}{(n+1)!n!} \right] \\&= \frac{1 \times 3 \times 5 \times \cdots \times (2n-1) \times 2 \times 4 \times 6 \times \cdots \times (2n)}{(n+1)!n!} \\&= \frac{(2n)!}{(n+1)n!n!} = \frac{1}{n+1} \binom{2n}{n}.\end{aligned}$$