

# Solving Recurrences

*Debdeep Mukhopadhyay*  
*IIT Kharagpur*

## Recurrence Relations

- A *recurrence relation* (R.R., or just *recurrence*) for a sequence  $\{a_n\}$  is an equation that expresses  $a_n$  in terms of one or more previous elements  $a_0, \dots, a_{n-1}$  of the sequence, for all  $n \geq n_0$ .
  - I.e., just a recursive definition, without the base cases.
- A particular sequence (described non-*recursively*) is said to *solve* the given recurrence relation if it is consistent with the definition of the recurrence.
  - A given recurrence relation may have many solutions.

## Example

- Consider the recurrence relation

$$a_n = 2a_{n-1} - a_{n-2} \quad (n \geq 2).$$

- Which of the following are solutions?

$$a_n = 3n$$

$$a_n = 2^n$$

$$a_n = 5$$

## Further Examples

- Recurrence relation for growth of a bank account with  $P\%$  interest per given period:

$$M_n = M_{n-1} + (P/100)M_{n-1}$$

- Growth of a population in which each pair of rabbit yield 1 new one every year after 2 years of their birth.

$$P_n = P_{n-1} + P_{n-2} \quad (\text{Rabbits and Fibonacci relation})$$

## Solving Compound Interest RR


- $$\begin{aligned}M_n &= M_{n-1} + (P/100)M_{n-1} \\ &= (1 + P/100) M_{n-1} \\ &= r M_{n-1} \quad (\text{let } r = 1 + P/100) \\ &= r (r M_{n-2}) \\ &= r \cdot r \cdot (r M_{n-3}) \quad \dots \text{and so on to } \dots \\ &= r^n M_0\end{aligned}$$

## Rabbits on an Island (assuming rabbits are immortal)

Year	Reproducing pairs	Young pairs	Total pairs
1	0	1	1
2	0	1	1
3	1	1	2
4	1	2	3
5	2	3	5
6	3	5	8


$$P_n = P_{n-1} + P_{n-2}$$

## Further Explanation



## Tower of Hanoi Example

- Problem: Get all disks from peg 1 to peg 2.
  - Only move 1 disk at a time.
  - Never set a larger disk on a smaller one.



*Question: Compute the number of steps  $H_n$*


Peg #1

Peg #2


Peg #3

# Intuition





$H_1=1$  is evident






So,  $H_2=3$



## For $n=3$ , $H_3=7$



1 step



## Hanoi Recurrence Relation

- Let  $H_n = \#$  moves for a stack of  $n$  disks.
- Optimal strategy:
  - Move top  $n-1$  disks to spare peg. ( $H_{n-1}$  moves)
  - Move bottom disk. (1 move)
  - Move top  $n-1$  to bottom disk. ( $H_{n-1}$  moves)
- Note:  $H_n = 2H_{n-1} + 1$

## Solving Tower of Hanoi RR

$$\begin{aligned}H_n &= 2 H_{n-1} + 1 \\&= 2 (2 H_{n-2} + 1) + 1 = 2^2 H_{n-2} + 2 + 1 \\&= 2^2 (2 H_{n-3} + 1) + 2 + 1 = 2^3 H_{n-3} + 2^2 + 2 + 1 \\&\dots \\&= 2^{n-1} H_1 + 2^{n-2} + \dots + 2 + 1 \\&= 2^{n-1} + 2^{n-2} + \dots + 2 + 1 \quad (\text{since } H_1 = 1) \\&= \sum_{i=0}^{n-1} 2^i \\&= 2^n - 1\end{aligned}$$

## Another R.R. Example

- Find a R.R. & initial conditions for the number of bit strings of length  $n$  without two consecutive 0s. Assume  $n \geq 3$ .
- We can solve this by breaking down the strings to be counted into cases that end in 0 and in 1.
  - For each ending in 0, the previous bit must be 1, and before that comes any qualifying string of length  $n-2$ .
  - For each string ending in 1, it starts with a qualifying string of length  $n-1$ .
- Thus,  $a_n = a_{n-1} + a_{n-2}$ . (Fibonacci recurrence.)

## Yet another R.R. example...

- Give a recurrence (and base cases) for the number of  $n$ -digit decimal strings containing an *even* number of 0 digits.
- Can break down into the following cases:
  - Any valid string of length  $n-1$  digits, with any digit 1-9 appended.
  - Any invalid string of length  $n-1$  digits, + a 0.
- $a_n = 9a_{n-1} + (10^{n-1} - a_{n-1})$   
 $= 8a_{n-1} + 10^{n-1}$ .
  - Base cases:  $a_0 = 1$  ( $\epsilon$ ),  $a_1 = 9$  (1-9).

# Catalan Numbers

- Find a R.R for the number of ways we can parenthesize the product of  $n+1$  numbers,  $x_0, x_1, \dots, x_n$  to specify the order of multiplication. Call it  $C_n$ .
- **Define  $C_0=C_1=1$  (its important to have proper base cases)**
- If  $n=2$ ,  $(x_0 \cdot x_1) \cdot x_2, x_0 \cdot (x_1 \cdot x_2) \Rightarrow C_2=2$ 
  - Note that  $C_2=C_1C_0+C_0C_1=1+1=2$
- If  $n=3$ ,  $((x_0 \cdot x_1) \cdot x_2) \cdot x_3; (x_0 \cdot x_1) \cdot (x_2 \cdot x_3); (x_0 \cdot (x_1 \cdot x_2)) \cdot x_3; x_0 \cdot ((x_1 \cdot x_2) \cdot x_3); x_0 \cdot (x_1 \cdot (x_2 \cdot x_3)) \Rightarrow C_3=5$ 
  - Note that  $C_3=C_2C_0+C_1C_1+C_0C_2=2+1+2=5$

# Catalan Numbers

- The final “.” operator is outside the scope of any parenthesis.
- The final . can be between any  $x_k$  and  $x_{k+1}$  out of the  $n+1$  numbers.
- How many ways can we have parenthesis as follows:

$$- [x_0, x_1, \dots, x_k] \cdot [x_{k+1}, x_{k+2}, \dots, x_n]$$

$C_k$

$C_{n-k-1}$

- The “.” can occur in after any  $x_k$ , where  $k$  ranges from 0 to  $n-1$
- So, the total number of possible parenthesis is:

$$\sum_{i=0}^{n-1} C_k C_{n-k-1}$$

**Exact form of  $C_n$   
can be computed using  
Generating functions**



## Solving Recurrences

- A linear homogeneous recurrence of degree  $k$  with constant coefficients (“ $k$ -LiHoReCoCo”) is a recurrence of the form

$$a_n = c_1 a_{n-1} + \dots + c_k a_{n-k},$$

where the  $c_i$  are all real, and  $c_k \neq 0$ .

- The solution is uniquely determined if  $k$  initial conditions  $a_0 \dots a_{k-1}$  are provided. This follows from the second principle of Mathematical Induction.

## Examples

- $f_n = f_{n-1} + f_{n-2}$  is a  $k$ -LiHoReCoCo
- $h_n = 2h_{n-1} + 1$  is not Homogenous
- $a_n = a_{n-1} + a_{n-2}^2$  is not linear
- $b_n = nb_{n-1}$  does not have a constant co-efficient

## Solving LiHoReCoCos

- The basic idea: Look for solutions of the form  $a_n = r^n$ , where  $r$  is a constant not zero ( $r=0$  is trivial)
- This requires the *characteristic equation*:  

$$r^n = c_1 r^{n-1} + \dots + c_k r^{n-k}, \text{ i.e., } \begin{matrix} \text{(rearrange} \\ \text{\& \times by } r^{k-n}) \end{matrix}$$

$$r^k - c_1 r^{k-1} - \dots - c_k = 0$$
- The solutions (*characteristic roots*) can yield an explicit formula for the sequence.

## Solving 2-LiHoReCoCos

- Consider an arbitrary 2-LiHoReCoCo:  

$$a_n = c_1 a_{n-1} + c_2 a_{n-2}$$
- It has the characteristic equation (C.E.):  

$$r^2 - c_1 r - c_2 = 0$$
- **Theorem 1**: If the CE has 2 roots  $r_1 \neq r_2$ , then  $\{a_n\}$  is a solution to the RR iff  $a_n = \alpha_1 r_1^n + \alpha_2 r_2^n$  for  $n \geq 0$  for constants  $\alpha_1, \alpha_2$ .

## Example

- Solve the recurrence  $a_n = a_{n-1} + 2a_{n-2}$  given the initial conditions  $a_0 = 2, a_1 = 7$ .
- Solution: Use theorem 1:
  - $c_1 = 1, c_2 = 2$
  - Characteristic equation:
 
$$r^2 - r - 2 = 0$$
  - Solutions:  $r = [ -(-1) \pm ((-1)^2 - 4 \cdot 1 \cdot (-2))^{1/2} ] / 2 \cdot 1$   
 $= (1 \pm 9^{1/2})/2 = (1 \pm 3)/2$ , so  $r = 2$  or  $r = -1$ .
  - So  $a_n = \alpha_1 2^n + \alpha_2 (-1)^n$ .

(Using the quadratic formula here.)

$$ax^2 + bx + c = 0 \Leftrightarrow$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

## Example Continued...

- To find  $\alpha_1$  and  $\alpha_2$ , solve the equations for the initial conditions  $a_0$  and  $a_1$ :
 
$$a_0 = 2 = \alpha_1 2^0 + \alpha_2 (-1)^0$$

$$a_1 = 7 = \alpha_1 2^1 + \alpha_2 (-1)^1$$
- Simplifying, we have the pair of equations:
 
$$2 = \alpha_1 + \alpha_2$$

$$7 = 2\alpha_1 - \alpha_2$$
- which we can solve easily by substitution:
 
$$\alpha_2 = 2 - \alpha_1; \quad 7 = 2\alpha_1 - (2 - \alpha_1) = 3\alpha_1 - 2;$$

$$9 = 3\alpha_1; \quad \alpha_1 = 3; \quad \alpha_2 = 1.$$
- Final answer:  $a_n = 3 \cdot 2^n - (-1)^n$

Check:  $\{a_{n \geq 0}\} = 2, 7, 11, 25, 47, 97 \dots$

## Proof of Theorem 1

- Proof that  $a_n = \alpha_1 r_1^n + \alpha_2 r_2^n$  is always a solution:
    - We know  $r_1^2 = c_1 r_1 + c_2$  and  $r_2^2 = c_1 r_2 + c_2$ .
    - Now we can show the proposed sequence satisfies the recurrence  $a_n = c_1 a_{n-1} + c_2 a_{n-2}$ :
 
$$\begin{aligned}
 c_1 a_{n-1} + c_2 a_{n-2} &= c_1(\alpha_1 r_1^{n-1} + \alpha_2 r_2^{n-1}) + c_2(\alpha_1 r_1^{n-2} + \alpha_2 r_2^{n-2}) \\
 &= \alpha_1 r_1^{n-2}(c_1 r_1 + c_2) + \alpha_2 r_2^{n-2}(c_1 r_2 + c_2) \\
 &= \alpha_1 r_1^{n-2} r_1^2 + \alpha_2 r_2^{n-2} r_2^2 = \alpha_1 r_1^n + \alpha_2 r_2^n = a_n. \quad \square
 \end{aligned}$$
- This shows that if  $a_n = \alpha_1 r_1^n + \alpha_2 r_2^n$ , then  $\{a_n\}$  is a solution to the R.R.**

## The remaining part of the proof

- If  $\{a_n\}$  is a solution of R.R. then,  $a_n = \alpha_1 r_1^n + \alpha_2 r_2^n$ , for  $n=0,1,2,\dots$
- Can complete proof by showing that for any initial conditions, we can find corresponding  $\alpha$ 's
  - $a_0 = C_0 = \alpha_1 + \alpha_2$
  - $a_1 = C_1 = \alpha_1 r_1 + \alpha_2 r_2$
  - $\alpha_1 = (C_1 - C_0 r_2) / (r_1 - r_2)$ ;  $\alpha_2 = (C_0 r_1 - C_1) / (r_1 - r_2)$
  - But it turns out this is a solution only if  $r_1 \neq r_2$ . So the roots have to be distinct.
  - The recurrence relation and the initial conditions determine the sequence uniquely. It follows that  $a_n = \alpha_1 r_1^n + \alpha_2 r_2^n$  (as we have already shown that this is a soln.)

## The Case of Degenerate Roots

- Now, what if the C.E.

$$r^2 - c_1r - c_2 = 0 \text{ has only 1 root } r_0?$$

- **Theorem 2:** Then,

$$\mathbf{a_n = (\alpha_1 + \alpha_2 n)r_0^n}, \text{ for all } n \geq 0,$$

for constants  $\alpha_1, \alpha_2$ .

## Example

- Solve:  $a_n = 6a_{n-1} - 9a_{n-2}$  with  $a_0 = 1, a_1 = 6$
- CE is  $r^2 - 6r + 9 = 0 \Rightarrow r_0 = 3$
- So, the general form of the soln is:
  - $a_n = (\alpha_1 + \alpha_2 n)3^n$
  - Solve the rest using the initial conditions

## ***k*-LiHoReCoCos**

- Consider a *k*-LiHoReCoCo:  $a_n = \sum_{i=1}^k c_i a_{n-i}$
- It's C.E. is:

$$r^k - \sum_{i=1}^k c_i r^{k-i} = 0$$

- **Thm.3:** If this has *k* distinct roots  $r_j$ , then the solutions to the recurrence are of the form:

$$a_n = \sum_{i=1}^k \alpha_i r_i^n$$

for all  $n \geq 0$ , where the  $\alpha_j$  are constants.

## **Example**

- Solve:

$a_n = 6a_{n-1} - 11a_{n-2} + 6a_{n-3}$ , with initial conditions  $a_0 = 2$ ,  $a_1 = 5$  and  $a_2 = 15$ .

CE is  $r^3 - 6r^2 + 11r - 6 = 0 \Rightarrow (r-1)(r-2)(r-3) = 0$

Thus the soln is:

$$a_n = (\alpha_1 1^n + \alpha_2 2^n + \alpha_3 3^n)$$

Solve the rest.

## Degenerate $k$ -LiHoReCoCos

- Suppose there are  $t$  roots  $r_1, \dots, r_t$  with multiplicities  $m_1, \dots, m_t$ . Then:

$$a_n = \sum_{i=1}^t \left( \sum_{j=0}^{m_i-1} \alpha_{i,j} n^j \right) r_i^n$$

for all  $n \geq 0$ , where all the  $\alpha$  are constants.

## Example

- Solve:  $a_n = -3a_{n-1} - 3a_{n-2} - a_{n-3}$ ,  $a_0 = 1$ ,  $a_1 = -1$ ,  $a_2 = -1$
- CE is :  $r^2 + 3r + 3r + 1 = (r+1)^3 = 0 \Rightarrow r = -1$  with multiplicity 3.
- So, soln is :
  - $a_n = (\alpha_1 + \alpha_2 r + \alpha_3 r^2)(-1)^n$
  - Complete the rest.

## LiNoReCoCos

- Linear *nonhomogeneous* RRs with constant coefficients may (unlike LiHoReCoCos) contain some terms  $F(n)$  that depend *only* on  $n$  (and *not* on any  $a_i$ 's). General form:

$$a_n = \underbrace{c_1 a_{n-1} + \dots + c_k a_{n-k}} + F(n)$$

The associated homogeneous recurrence relation (associated LiHoReCoCo).  $\mathbf{F(n)}$  is **not identically zero**.

## Solutions of LiNoReCoCos

- A useful theorem about LiNoReCoCos:
  - If  $a_n = p(n)$  is any *particular* solution to the LiNoReCoCo

$$a_n = \left( \sum_{i=1}^k c_i a_{n-i} \right) + F(n)$$

- Then *all* its solutions are of the form:

$$a_n = p(n) + h(n),$$

where  $a_n = h(n)$  is any solution to the associated homogeneous RR  $a_n = \left( \sum_{i=1}^k c_i a_{n-i} \right)$



## LiNoReCoCo Example

- Find all solutions to  $a_n = 3a_{n-1} + 2n$ . Which solution has  $a_1 = 3$ ?
  - Notice this is a 1-LiNoReCoCo. Its associated 1-LiHoReCoCo is  $a_n = 3a_{n-1}$ , whose solutions are all of the form  $a_n = \alpha 3^n$ . Thus the solutions to the original problem are all of the form  $a_n = p(n) + \alpha 3^n$ . So, all we need to do is find one  $p(n)$  that works.

## Trial Solutions

- If the extra terms  $F(n)$  are a degree- $t$  polynomial in  $n$ , you should try a degree- $t$  polynomial as the particular solution  $p(n)$ .
- This case:  $F(n)$  is linear so try  $a_n = cn + d$ .
$$cn + d = 3(c(n-1) + d) + 2n \quad (\text{for all } n)$$
$$(2c+2)n + (2d-3c) = 0 \quad (\text{collect terms})$$

So  $c = -1$  and  $d = -3/2$ .

So  $a_n = -n - 3/2$  is a solution.
- Check:  $a_{n \geq 1} = \{-5/2, -7/2, -9/2, \dots\}$

## Finding a Desired Solution

- From the previous, we know that all general solutions to our example are of the form:

$$a_n = -n - 3/2 + \alpha 3^n.$$

Solve this for  $\alpha$  for the given case,  $a_1 = 3$ :

$$3 = -1 - 3/2 + \alpha 3^1$$

$$\alpha = 11/6$$

- The answer is  $a_n = -n - 3/2 + (11/6)3^n$ .

## Double Check Your Answer!

- Check the base case,  $a_1=3$ :

$$a_n = -n - 3/2 + (11/6)3^n$$

$$a_1 = -1 - 3/2 + (11/6)3^1$$

$$= -2/2 - 3/2 + 11/2 = -5/2 + 11/2 = 6/2 = 3$$

- Check the recurrence,  $a_n = 3a_{n-1} + 2n$ :

$$-n - 3/2 + (11/6)3^n = 3[-(n-1) - 3/2 + (11/6)3^{n-1}] + 2n$$

$$= 3[-n - 1/2 + (11/6)3^{n-1}] + 2n$$

$$= -3n - 3/2 + (11/6)3^n + 2n = -n - 3/2 + (11/6)3^n \blacksquare$$

## Theorem

- Suppose that  $\{a_n\}$  satisfies the LiNoReCoCo,
 
$$a_n = c_1 a_{n-1} + \dots + c_k a_{n-k} + F(n),$$
 where  $c_1, c_2, \dots, c_k$  are real numbers and  $F(n) = (b_t n^t + b_{t-1} n^{t-1} + \dots + b_0) s^n$ , where  $b$ 's and  $s$  are real numbers.
- When  $s$  is not a root of the CE, there is a Particular solution of the form:  $(p_t n^t + p_{t-1} n^{t-1} + \dots + p_0) s^n$ .
- When  $s$  is a root of this CE and its multiplicity is  $m$ , there is a particular solution of the form:  $n^m (p_t n^t + p_{t-1} n^{t-1} + \dots + p_0) s^n$

## State the Particular Solutions

- RR:  $a_n = 6a_{n-1} - 9a_{n-2} + F(n)$
- CE has a single root 3, with multiplicity 2.
- $F(n) = 3^n$  Particular Solution:  $p_0 n^2 3^n$
- $F(n) = n 3^n$  Particular Solution:  $n^2 (p_1 n + p_0) 3^n$
- $F(n) = n^2 2^n$  Particular Solution:  $(p_2 n^2 + p_1 n + p_0) 2^n$
- $F(n) = (n^2 + 1) 3^n$  Particular Solution:  $n^2 (p_2 n^2 + p_1 n + p_0) 3^n$

## Be Careful when $s=1$

- Example:  $a_n = a_{n-1} + n$ ,  $a_1 = 1$
- CE:  $r=1$ , with multiplicity 1
- $F(n)=n$ , Particular Solution is  $n(p_1n+p_0)$
- Solve for  $p_1$  and  $p_0$  using the recurrence equation
- Write the solution:  $c$  (solution to the associated homogenous RR) + Particular Solution
- Solve for  $c$  using  $a_1=1$  and obtain  $a_n = n(n+1)/2$